

Anti-Rb phospho-Ser795 antibody, monoclonal (28B5)

71-171 100 ug

Retinoblastoma protein (Rb), the tumor suppressor product of the retinoblastoma susceptibility gene, is a 110 kDa protein that functions as a negative regulator of the cell cycle by arresting cells in the G1 phase and halting inappropriate cell proliferation (1). At the transcriptional level, Rb protein exerts its growth suppressive function by binding to transcription factors including E2F1, PU1, ATF2, UBF, Elf-1 and c-Abl (2-4). Loss of Rb function leads to uncontrolled cell growth and tumor development and is found in all retinoblastomas and in a variety of other human malignancies. The ability of Rb protein to inhinbit transcription and cell cycle progression is inactivated by phosphorylation, which is catalyzed by the cyclin-dependent protein kinases.

Applications

1. Western blotting (~1ug/ml)

2. ELISA

Other applications were not tested.

Specification

Antigen: A synthetic peptide containing phospho-Ser795 of human Rb

Isotype: Mouse IgG2a (κ)

Form: Purified monoclonal antibody (IgG) 1mg/ml in PBS, 50% glycerol, filter-sterilized

Specificity: Specific to human Rb phosphorylated at Ser795. Not tested for other species.

Can detect endogenous levels of Rb phosphorylated at Ser795 in most cell lines.

Storage: -20 °C (long period, -70°C)

Data Link UniProtKB/Swiss-Prot P06400 (RB_HUMAN)

References

- 1. Sherr CJ "Cancer cell cycles" Science 274: 1672-1677 (1996) PMID: 8939849
- 2. Nevins JR "E2F: a link between the Rb tumor suppressor protein and viral oncoproteins" *Science* **258**: 424-429 (1992) PMID: <u>1411535</u>
- 3. Welch PJ & Wang JY "A C-terminal protein-binding domain in the retinoblastoma protein regulates nuclear c-AbI tyrosine kinase in the cell cycle" *Cell* 75: 779-790 (1993) PMID: 8242749
- 4. Hu QJ et al "The regions of the retinoblastoma protein needed for binding to adenovirus E1A or SV40 large T antigen are common sites for mutations" EMBO J 9:1147-1155 (1990) PMID: 2138977

Fig.1 Specificity of the monoclonal antibody (28B5) to the phosphorylated Rb at Ser795 as demonstrated by Western blotting.

Crude extracts of human lung carcinoma cell line H1299 transfected with plasmid expressing Myc-tagged wild-type Rb (lane 1), Rb (S795A) (lane 2) or Rb (T821A) were immuno-precipitated with anti-Myc anti-body and the precipitates were analyzed by Western blotting with monoclonal antibody 28B5.(Data provided by Dr. Y. Inoue at the JFCR Cancer Research Institute, Tokyo)

Related product: #71-173 anti-Rb phospho-Thr821 antibody, monoclonal (24A7)

BioAcademia, Inc. Tel. 81-6-6877-2335 Fax. 81-6-6877-2336 info@bioacademia.co.jp http://www.bioacademia.co.jp/en/