

Anti-mouse ADAM1B antibody, rat monoclonal (#57)

73-010 100 μg

Storage: Shipped at 4° C and store at -20° C...

Reactivity: Mouse.

Applications:

1.Immunocytochemistry (1/100 - 1/300)

2. Immunohistochemistry (1/100~1/300) paraffin-embedded section.

Not suitable for western blotting.

Recommend to use clone #158 (BioAcademia 73-007) for western blotting.

Immunogen: Mouse sperm

Form: Purified monoclonal antibody (IgG) 1mg/ml in PBS, 50% glycerol, filter-sterilized.

Azide- and carrier-free.

Key words: Acrosome reaction, Membrane fusion, Protein trafficking, IZUMO1,

Sperm-egg fusion

Function: May play a role in spermatogenesis, sperm maturation and fertilization...

Molecular mass: 89,369 with 806 amino acids. N-terminal signal peptide with 33 amino acids from this protein is processed to give propertide, which may undergo further processing.

Database Links: UniProtKB Q8R534 (mouse ADAM1b)

Reference: This antibody was described in and used in the following publications.

Ikawa M. et al. Calsperin is a testis-specific chaperone required for sperm fertility.

J Biol Chem. 2011;286:5639-46. IF

Satouh Y., et al (2012) Visualization of the moment of mouse sperm—egg fusion and dynamic localization of IZUMO1. J. Cell Science 125, 4985–4990. PubMed 22946049. **IF.**




Fig.1 Immunofluorescent staining of ADAM1B in sperm by using anti-ADAM1B antibody.

.Fresh sperm collected from the epididymis of mice were fixed in 4% paraformaldehyde and then immunostained with the anti-ADAM1B antibody (#57) at 1/300 dilution.

BioAcademia, Inc. Tel. 81-6-6877-2335 Fax. 81-6-6877-2336 info@bioacademia.co.jp http://www.bioacademia.co.jp/en/






Fig.2. Immunohistological staining of ADAM1B in mouse testis using anti-ADAM1B antibody (#57). A section of formalin fixed and paraffin embedded mouse testis was treated with the anti-ADAM1B antibody at 1/100 dilution after deparaffization and antigen retrieval. The 2nd antibody, anti-rat IgG conjugated with Alexa Fluor 488 (Abcam) was used at 1/1,000 dilution. DNA was stained with DAPI and the merged image was shown (Merge). The bright field microsopic picture of the same region was shown on the right.